

Fertility and Inheritance of Variegated and Purple Foliage Across a Polyploid Series in *Hypericum androsaemum* L.

Richard T. Olsen¹ and Thomas G. Ranney²

Department of Horticultural Science, Mountain Horticultural Crops Research and Extension Center, North Carolina State University, Fletcher, NC 28732-9244

Dennis J. Werner³

Department of Horticultural Science, North Carolina State University, Raleigh, NC 27695-7609

ADDITIONAL INDEX WORDS. st. john's wort, tutsan, autotetraploid, oryzalin, invasive plants, pollen viability, ornamental plant breeding

ABSTRACT. Inheritance of two mutant foliage types, variegated and purple, was investigated for diploid, triploid, and tetraploid tutsan (*Hypericum androsaemum*). The fertility of progeny was evaluated by pollen viability tests and reciprocal crosses with diploids, triploids, and tetraploids and germinative capacity of seeds from successful crosses. Segregation ratios were determined for diploid crosses in reciprocal di-hybrid F_1 , F_2 , BCP1, and BCP2 families and selfed F_2 s with the parental phenotypes. F_2 tetraploids were derived from induced autotetraploid F_1 s. Triploid segregation ratios were determined for crosses between tetraploid F_2 s and diploid F_1 s. Diploid di-hybrid crosses fit the expected 9:3:3:1 ratio for a single, simple recessive gene for both traits, with no evidence of linkage. A novel phenotype representing a combination of parental phenotypes was recovered. Data from backcrosses and selfing support the recessive model. Both traits behaved as expected at the triploid level; however, at the tetraploid level the number of variegated progeny increased, with segregation ratios falling between random chromosome and random chromatid assortment models. We propose the gene symbol *var* (variegated) and *pl* (purple leaf) for the variegated and purple genes, respectively. Triploid pollen stained moderately well (41%), but pollen germination was low (6%). Triploid plants were highly infertile, demonstrating extremely low male fertility and no measurable female fertility (no viable seed production). The present research demonstrates the feasibility of breeding simultaneously for ornamental traits and non-invasiveness.

The genus *Hypericum* L. is a member of the large family Guttiferae (Clusiaceae, subfamily Hypericoideae, tribe Hypericeae) (Gustafsson et al., 2002) and contains about 370 species of herbs, shrubs, and trees distributed across the temperate zones and high elevations of the tropics (Mabberley, 1997; Robson, 1985). St. john's worts, as they are collectively known, have garnered much attention of late, with research focused on phytochemicals unique to the genus, including hypericins and hyperforins from common st. john's wort (*H. perforatum* L.) (Kirakosyan et al., 2004) and xanthones from tutsan (*H. androsaemum*) (Dias et al., 2000; Valentão et al., 2002). In addition to pharmacological value, the genus has contributed a number of important ornamental species including *H. calycinum* L., *H. forrestii* (Chittenden) N. Robson, and *H. androsaemum* (Robson, 1985).

Hypericum androsaemum ($2n = 2x = 40$) is native from western Europe, through the Mediterranean and northern Africa, east into northern Iran (Robson, 1985). This semi-evergreen shrub grows less than 1 m in height and has an abundance of yellow, star-like flowers and fleshy red fruit that mature to a brown capsule. As a landscape plant, *H. androsaemum* and its hybrids with *H. hircinum* L. (*H. xinodorum* Miller), perform best in the cooler parts of U.S. Department of Agriculture hardiness zones 6 to 8. Various cultivars have been selected for the landscape and floriculture trade including yellow, variegated, and purple leaf forms. 'Mrs. Gladis Brabazon' and 'Glacier' are cultivars derived from *H. androsaemum* f. *variegatum* McClintock & Nelson with heavily mottled leaves consisting of white, green, and light green patches (McClintock et al., 1986). 'Albury Purple' has purple blushed foliage and red-veined flowers; the origin of this cultivar is unknown.

Cultivation of *H. androsaemum* outside of its native range has led to naturalization in Europe, Australia, New Zealand, and Chile (Robson, 1985) and is a species of concern in the Pacific northwestern U.S. (D. Hinkley, personal communication). It is considered an invasive species in Australia, where biological control mechanisms are being studied (McLaren et al., 1997). Increasing awareness and concern about invasive plant species (Reichard and White, 2001) provide an opportunity for plant breeders to address this problem through the development of non-invasive cultivars. Bananas (*Musa* L.) and watermelons [*Citrullus lanatus* (Thunb.) Matsum. & Nakai] are well-known examples of triploid plants displaying low fertility and seed development (Kihara, 1951; Ortiz and Vuylsteke, 1995). Few triploid ornamental plants have been developed, though the potential for using this approach to

Received for publication 26 Apr. 2006. Accepted for publication 28 Sept. 2006. This research was funded, in part, by the North Carolina Agricultural Research Service (NCARS), Raleigh, NC 27695-7643; the North Carolina Association of Nurserymen, Inc., 969 Trinity Road, Raleigh, NC 27607; U.S. Dept. of Agriculture Floral and Nursery Research Initiative, Beltsville, MD 20705-2350; and J. Frank Schmidt Family Charitable Foundation, Boring, OR 97009. Use of trade names in this publication does not imply endorsement by NCARS of the products named nor criticism of similar ones not mentioned. Technical assistance of Thomas Eaker, Joel Mowrey, and Nathan Lynch, and staff of the Mountain Horticultural Crops Research and Extension Center is greatly appreciated. From a dissertation submitted by R.T.O. in partial fulfillment of the requirements for the PhD degree.

¹Corresponding author. Current address: Floral and Nursery Plants Research Unit, USDA-ARS, US National Arboretum, 3501 New York Ave. NE, Washington, DC 20002; e-mail: Richard.Olsen@ars.usda.gov

²Professor.

³Professor, Director of the J.C. Raulston Arboretum.

develop non-invasive cultivars is promising (Ranney, 2004). The triploid rose-of-sharon (*Hibiscus syriacus* L.) cultivars released by the U.S. National Arboretum shrub breeding program are notable exceptions (Egolf, 1970, 1981, 1986, 1988).

The objectives of this study were to investigate the mode of inheritance for two important ornamental traits, leaf variegation and purple foliage, across three ploidy levels [diploid (2x), triploid (3x), and tetraploid (4x)] and to determine the fertility of progeny to assess the feasibility of developing non-invasive, triploid cultivars.

Materials and Methods

INHERITANCE STUDIES. Plants of *H. androsaemum* f. *variegatum* ‘Glacier’ (G) and ‘Albury Purple’ (AP) were maintained in the ornamental tree and shrub collection at the Mountain Horticultural Crops Research Station (MHCRC), Fletcher, N.C. Prior to beginning the study, open-pollinated seedlings from G and AP displayed their parental phenotypes, indicating that both traits were heritable. Plants for all crosses were grown in the greenhouse at MHCRC in 18.9-L containers in 3 pine bark : 1 peat (by volume) substrate amended with 2.8 kg·m⁻³ dolomitic limestone and 0.5 kg·m⁻³ micronutrients (Micromax; The Scotts Co., Marysville, Ohio), top-dressed with 50 g per container of 15N–3.9P–10.0K controlled-release fertilizer (15–9–12 Osmocote Plus 3–4 months at 70 °C; The Scott’s Co.) and watered by hand as needed. Reciprocal, di-hybrid F₁ and F₂ families were generated to investigate inheritance (Table 1). BCP1 and BCP2 families were produced as additional test crosses. F_{2P1(2x)} plants exhibiting the original parental phenotypes were selfed (S₀) to further verify genotypes. For all crosses, flowers were emasculated prior to anthesis, generally when the elongating floral bud was equal to subtending calyx lobes in length. Pollen was collected from recently dehiscent anthers and either used fresh or dried overnight at 5 °C using indicator drierite (Drierite, Xenia, Ohio) and stored at 5 °C for use in later crosses. Pollen was applied daily to stigmas using small brushes, until stigmas turned brown after ~10 d. Fruit were collected when capsules turned from bright red to brown and allowed to dry at room temperature for ~2 d. Seeds were surface-sown onto a substrate of 1 peat : 1 vermiculite (by volume) in seedling flats (58 × 36 × 6 cm) and watered with intermittent mist (8 s every 6 min for 24 h) until germination occurred in 1 to 4 weeks. Seedlings were transplanted into 40-cell trays with the same substrate and phenotypes were scored when seedlings had three sets of true leaves.

A subset of seedlings from G × AP F₁ [F_{1P1(2x)}] were treated with the mitotic-inhibiting herbicide oryzalin to induce tetraploidy

Table 1. Crosses between *Hypericum androsaemum* ‘Glacier’ (G) and ‘Albury Purple’ (AP) phenotypes and families produced.

Cross	Ploidy of progeny	Families
G × AP	2x	F _{1P1(2x)} , F _{2P1(2x)} ^z
AP × G	2x	F _{1P2(2x)} , F _{2P2(2x)}
G × F _{1P1(2x)}	2x	BCP1
AP × F _{1P1(2x)}	2x	BCP2
G F _{2P1(2x)} selfed	2x	S ₀
AP F _{2P1(2x)} selfed	2x	S ₀
F _{1P1(4x)} × F _{1P1(2x)}	3x	F _{2(3x)}
F _{1P1(4x)} selfed	4x	F _{2(4x)}

^zPloidy level in parentheses for each filial generation for clarity in describing subsequent crosses.

[F_{1P1(4x)}]. A total of 488 seedlings were selected from seedling flats after the first set of true leaves had emerged. Substrate was washed from roots and the entire seedling submerged in 150 μM oryzalin [0.004% Surflan; Dow AgroSciences, Indianapolis] for 24 h. Following treatment, seedlings were washed under running water for 30 min and potted into 40-cell trays containing a 1 peat : 1 vermiculite (by volume) substrate and placed under intermittent mist (8 s every 6 min for 24 h) for 3 d, then grown under normal greenhouse conditions. Relative DNA content and associated ploidy levels were determined using a flow cytometer (PA-I Ploidy Analyzer; Partec GmbH, Münster, Germany). Nuclei isolation and staining [4',6-diamidino-2-phenylindole (DAPI)] followed the protocols provided by (Olsen et al., 2006). Induced tetraploids [F_{1P1(4x)}] were grown to flowering and selfed to investigate segregation of traits in an F₂ tetraploid family [F_{2(4x)}] or pollinized with diploid F₁s (F_{1P1(2x)}) to investigate segregation at the triploid level [F_{2(3x)}] (Table 1). Samples of seedlings (n = 25) from all crosses were analyzed using flow cytometry to confirm ploidy level of the progeny. Flow cytometry was also used to detect apomixis by analyzing ploidy levels of seeds from F₁ crosses (Table 1). Seeds from sexual crosses will have a diploid embryo peak and a triploid endosperm peak, while seeds from apomictic (unreduced embryo sac) origin will have a diploid embryo peak and a tetraploid endosperm peak (Matzk et al., 2000). Seeds (n = 25 per sample) were chopped using the same buffer and staining procedure as for foliage.

We hypothesized that variegated and purple leaf traits were two independent loci, both inherited in a simple Mendelian recessive manner. Segregation data were analyzed for departures from expected ratios using chi-square analysis on segregating families [F_{2P1(2x)}, F_{2P2(2x)}, BCP1, BCP2, F_{2(3x)}, and F_{2(4x)}]. The chi-square test of independence for linkage was calculated for diploid F₂s [F_{2P1(2x)} and F_{2P2(2x)}]. Crosses were conducted during the summers of 2003–05.

FERTILITY OF PROGENY. Reciprocal crosses between diploid (2x), triploid (3x), and tetraploid plants (4x) were made in 2005 to investigate male and female fertility as a function of ploidy level. Plants were grown in 18.9-L containers in the greenhouse using the same substrate and conditions described earlier and the same pollination techniques. Six plants per ploidy level were randomly sampled for pollen viability during the study. One flower per plant per ploidy level was harvested and anthers divided into two groups, with one group for pollen staining and the other for pollen germination tests. Pollen grains were stained in 40 μL of 1% acetocarmine stain on a microscope slide. The slide was heated 3× over a hot-plate for 5 s, sealed with valap (1 vaseline : 1 lanolin : 1 paraffin by weight), and incubated at room temperature (23 °C) for 5 h. Stained, well-formed pollen grains were scored as viable. Pollen germination was performed in 5-mL petri dishes containing Brewbaker-Kwack media supplemented with 5% sucrose and solidified with 2% agarose (Marquard, 1992). Pollen grains with pollen tubes greater than one-half the diameter of the pollen grain after 5 h were scored as germinated. Pollen staining and germination were observed using a compound light microscope (Micromaster; Fisher Scientific, Pittsburgh) under ×100 and ×400 magnifications. For each replicate ≥100 pollen grains were scored. The experiment was a completely randomized design with three treatments (ploidy levels) and six replicates (n = 6).

Crosses among the three ploidy levels resulted in nine factorial treatment combinations. The experiment was a completely randomized design with five single-plant replicates (n = 5) per

cross with 10 flowers (subsamples) pollinated per plant. Fruit from successful crosses matured ~40 d after pollination. Fruit were harvested, dried, and seeds separated and stored in sealed vials at 5 °C until used for germination studies. Total number of fruit per cross, percentage of fruit set, and average seeds per fruit were recorded.

Germinative capacity of seeds harvested from successful crosses was determined using in vitro germination studies. Two germination blotters (SDB3.5; Anchor Paper Co., St. Paul, Minn.) soaked with 100 μ M gibberellic acid (GA₄₊₇) (21 g·L⁻¹ Provide; Valent Biosciences, Walnut Creek, Calif.) were placed in petri dishes (100 × 15 mm). A sample of 50 seeds per fruit, or all seeds available when <50, were sown per petri dish. Petri dishes were sealed with parafilm and placed in a growth chamber maintained at 22 °C, 16 h day/8 h night period, and 42 μ mol·s⁻¹·m⁻² photosynthetically active radiation (PAR) provided by fluorescent lights. After 8 weeks, seeds with emergent radicles were scored as germinated.

Data from pollen, crossing, and germination studies were analyzed using analysis of variance (PROC GLM; SAS version 8.02, SAS Institute Inc., Cary, N.C.) and means compared using Fisher's protected least significant difference (LSD) at $\alpha = 0.05$.

Results

INHERITANCE STUDIES. Reciprocal F₁ crosses yielded progeny that were all non-variegated and green, with the exception of three variegated seedlings [F_{1P1(2x)}] and one purple seedling [F_{1P2(2x)}], which were attributed to accidental self-pollination (Table 2). No evidence for apomixis was observed from flow cytometry analysis of F₁ seed; thus, all progeny appeared to be the result of sexual crosses (data not shown). Segregation ratios for the diploid F₂ families [F_{2P1(2x)} and F_{2P2(2x)}] fit the expected 9:3:3:1 ratio, supporting the hypothesis of simple recessive inheritance for both traits. The chi-square test for independence of linkage [$\chi^2 = 1.75$, $P = 0.19$ and $\chi^2 = 0.02$, $P = 0.89$ for F_{2P1(2x)} and F_{2P2(2x)}, respectively] suggests no evidence for linkage between the variegated and purple loci. Furthermore, recovery of variegated and purple seedlings in both F₂ populations [F_{2P1(2x)} and F_{2P2(2x)}] effectively

rules out strict maternal inheritance. Segregation ratios for BCP1 fit the expected 1:1 ratio of non-variegated:variegated phenotypes (Table 2). However, in BCP2 a few spontaneous variegated green and variegated purple seedlings were observed which can not be explained by accidental self-pollination or pollen contamination. These seedlings were scored as either non-variegated or purple for chi-square analysis; however, due to the presence of more purple seedlings than expected (221 observed vs. 199.5 expected) we still had a poor fit. F₂ plants [F_{2P1(2x)}] exhibiting the original G or AP phenotype when selfed (S₀) yielded all variegated or all purple progeny, respectively (Table 2).

Induction of autotetraploids [F_{1P1(4x)}] from F_{1P1(2x)} seedlings was successful. Of 488 seedlings treated, 16.2% survived (79 of 488) the treatment. Of those that survived, 39.2% remained diploid, 45.6% were mixoploids, and 15.2% were non-chimeric tetraploids. Autotetraploids grew slowly at first, but otherwise were similar to their diploid counterparts. Triploid [F_{2(3x)}] and tetraploid [F_{2(4x)}] seedlings from crosses with autotetraploids grew normally, except tetraploid AP phenotypes which tended to be poor growers.

Triploid F₂ [F_{2(3x)}] progeny fit the expected 121:11:11:1 segregation ratio (Table 2). However, at the tetraploid level [F_{2(4x)}] we observed more than three times as many variegated phenotypes than expected (76 observed vs. 21.6 expected) and thus had a poor fit to the expected 1225:35:35:1 segregation ratio for tetrasomic inheritance at the tetraploid level. We analyzed inheritance for each locus separately for random chromosome and chromatid assortment (at 50% recombination), as well as for selective pairing at the tetraploid level to elucidate possible mechanisms for departure from our expected ratio for the variegated trait. Random chromosome assortment for one recessive gene at the tetraploid level would segregate 35 wild-type:1 mutant. For the purple locus, there was no reason to reject the assumption of random chromosome assortment at the tetraploid level ($\chi^2 = 0.028$, $P = 0.867$); however, the variegated locus did not fit ($\chi^2 = 138.9$, $P < 0.001$). For random chromatid assortment at 50% recombination we would expect 20.8 wild-type:1 mutant, and the variegated locus failed to fit ($\chi^2 = 46.4$, $P < 0.001$). Assuming preferential or selective pairing, we would expect 15 wild-type:1 mutant, and

Table 2. Segregation for foliage traits in *Hypericum androsaemum* across ploidy levels in families derived from 'Glacier' (G) and 'Albury Purple' (AP).

Cross	Families	Progeny (no. seedlings) ^z				Expected ratio	χ^2	P
		+/-	var/var	+/-	pl/pl			
G x AP	F _{1P1(2x)} ^y	130	3 ^x	0	0	1:0:0:0	0.00	1.00
AP x G	F _{1P2(2x)}	162	0	1 ^x	0	1:0:0:0	0.00	1.00
G x AP	F _{2P1(2x)}	218	71	88	20	9:3:3:1	3.69	0.30
AP x G	F _{2P2(2x)}	214	77	81	28	9:3:3:1	1.43	0.70
G x F _{1P1(2x)}	BCP1	200	196	0	0	1:1:0:0	0.04	0.84
AP x F _{1P1(2x)}	BCP2	178	4 ^w	221	3 ^w	1:0:1:0	4.63	0.03
G selfed	S ₀	0	200	0	0	0:1:0:0	0.00	1.00
AP selfed	S ₀	0	0	200	0	0:0:1:0	0.00	1.00
F _{1P1(4x)} x F _{1P1(2x)}	F _{2(3x)}	573	46	49	3	121:11:11:1	1.38	0.71
F _{1P1(4x)} selfed	F _{2(4x)}	701	76	22	1	1225:35:35:1	141.20	<0.001

^zNumber of progeny for each phenotypic class (+/- +/- = non-variegated, green; var/var +/- = variegated, green; +/- pl/pl = non-variegated, purple; var/var pl/pl = variegated, purple).

^yPloidy level in parentheses for each filial generation for clarity in describing subsequent crosses.

^xUnexpected and inconsistent with proposed simple recessive model, due to possible accidental self-pollination. Data not used in chi-square calculations.

^wUnexpected and unexplained variegation, which can not be explained by self- or cross-pollen contamination. These seedlings were scored as either non-variegated or purple for chi-square calculations.

again, the variegated locus failed to fit ($\chi^2 = 15.6$, $P < 0.001$).

If above the diploid level the variegated trait behaves quantitatively, the frequency of variegated progeny in both triploid and tetraploid crosses would increase, and classes of variegation from weak (duplex) to strong variegation (quadriplex) may be discerned. At the tetraploid level we would then expect 3 variegated : 1 non-variegated progeny; however, the variegated locus did not fit this either ($\chi^2 = 100.9$, $P < 0.001$).

FERTILITY OF PROGENY. Ploidy level had a significant effect on pollen staining (F value = 274.9, $P < 0.0001$) and germination (F value = 136.3, $P < 0.0001$). Diploid (2x) and tetraploid (4x) plants had greater pollen staining (96.8% and 94.1%, respectively) and germination (89.1% and 76.3%, respectively) than triploids (3x) (40.6% staining and 5.9% germination).

Crosses between diploids (2x) resulted in 64.0% fruit set with an average of 632 seeds per fruit and a germination percentage of 26.3% (Table 3). Only two fruit formed in 2x x 3x crosses, with an average of 138 seeds per fruit which germinated at 1.0%. Crosses between 2x x 4x were not successful. Triploids failed to set fruit and viable seeds when pollinated with 2x or 3x pollen. One fruit was collected from 3x x 4x crosses, yielding four seeds which failed to germinate. Tetraploids were successfully crossed with 2x and 4x pollen, resulting in similar percentage fruit sets, average seeds per fruits and seed germination values. Crosses between 4x x 3x were unsuccessful. Germination was erratic for diploid (2x x 2x), triploid (4x x 2x) and tetraploid (4x x 4x) seed, as evident in the very broad least significant difference (LSD_{0.05}) for seed germination (28.4%).

Discussion

Reciprocal di-hybrid crosses were performed to rule out maternal effects on inheritance of each trait, in particular, variegation. Variegated foliage is often the result of abnormal plastid development due to mutations in chloroplast genes and can result in strict maternal inheritance or non-Mendelian inheritance due to stochastic sorting of aberrant vs. normal plastids during cell division and transmission to germ cells (Tilney-Bassett, 1978; Walbot and Coe, 1979). F₁ reciprocal crosses of G x AP resulted in non-variegated, green progeny (wild-type), indicating that

variegation in *H. androsaemum* f. *variegatum* 'Glacier' is not the result of maternally (or paternally) transmitted aberrant plastids. Recovery of variegated progeny in F₂s at the expected 9:3:3:1 segregation ratio and in BCP1 at 1:1 is evidence that variegation is controlled by a monogenic, recessive nuclear gene. We propose the gene symbol *var* (variegated) for the recessive allele producing the variegated phenotype (*var/var*) of *H. androsaemum* f. *variegatum* and the cultivars Mrs. Gladis Brabazon and Glacier derived from this forma. Presence of variegation only in homozygous genotypes, with no aberrant plastid transmission, suggests the variegated allele is a mutation at a locus that regulates chloroplast gene expression or encodes a protein necessary for proper functioning of the chloroplast. Plastid restitution occurs when at least one wild-type allele is present, as in the *immutans* mutant of *Arabidopsis thaliana* (L.) Heynh. (Aluru and Roderme, 2004).

The lack of purple foliage in F₁s and recovery of purple progeny at the expected ratio in the F₂s supports our hypothesis of simple recessive inheritance for purple foliage. We propose the gene symbol *pl* (purple leaf) for the recessive allele originating from *H. androsaemum* 'Albury Purple.' Purple (or red) foliage variants, resulting from foliar accumulation of anthocyanins, have variously been described in agronomic crops and are important in ornamental plant breeding programs. Cadic (1992) reported inheritance of purple foliage in barberries (*Berberis* L.) due to a single recessive gene, as we observed in *H. androsaemum*. In peach [*Prunus persica* (L.) Batsch.] red foliage is controlled by a single locus, *Gr*, and with the *Gr* allele incompletely dominant (Chaparro et al., 1995), as is bronze-leaf (*Rt*) in *Malus* Mill. (Alston et al., 2000; Sampson and Cameron, 1965). The lack of purple pigmentation in our F₁s and no variation in purple foliage intensity in our segregating progeny suggests totally recessive gene action. In other woody plant genera, inheritance of purple foliage is dominant as in *Betula pendula* Roth. 'Purpurea' (Hattemer et al., 1990), *Fagus sylvatica* L. (Blinkenberg et al., 1958; Heinze and Geburek, 1995), and *Corylus* L. (Smith and Mehlenbacher, 1996; Thompson, 1985), though variation in coloration exists in *Corylus* due to different alleles and complimentary gene action. Backcrossing to AP(BCP2) resulted in the recovery of more purple seedlings than expected; however, these may be explained by accidental self-pollination. Recovery of a few spontaneous variegated progeny was not expected in BCP2, and not explained by selfing or pollen contamination. Phenotypes were scored at the three-leaf stage, and these seedlings may represent the accumulation of deleterious alleles unrelated to the variegated locus or instability of our variegated locus due to a transposable element.

Recovery in F₂ lines of individuals with both variegated and purple foliage at the expected ratio indicates that the two traits (loci) are not closely linked. Based on a simple recessive model of inheritance for both variegated and purple foliage, phenotypes displaying the novel combination of variegation on purple foliage have the genotype *var/var pl/pl*.

Both traits behaved as expected at the triploid level, but at the tetraploid level we recovered more variegated, green progeny than expected. At the tetraploid level, segregation ratios for the variegated trait fell between that expected with random chromosome assortment and random chromatid assortment for autotetraploids. Assuming random pairing between homologous chromosomes and independent assortment, we would expect perfect tetrasomic inheritance and 35 non-variegated : 1 variegated progeny in the F_{2(4x)}. However, the prevalence of multivalent formation and frequency of recombination will influence double reduction and

Table 3. Fertility as a function of ploidy in *Hypericum androsaemum* reciprocal interpollid crosses.

Female parent	Male parent	Fruit developed			Seed germination (%) ^y
		(no.)	(%) ^z	(avg no. seeds) ^y	
2x	2x	32	64.0 a ^x	632 a	26.3 a
2x	3x	2	2.0 b	138 ab	1.0 a
2x	4x	0	0.0	---	---
3x	2x	0	0.0	---	---
3x	3x	0	0.0	---	---
3x	4x	1	4.2 b	4.0 b	0.0 a
4x	2x	7	10.8 b	260 ab	5.7 a
4x	3x	0	0.0	---	---
4x	4x	15	20.7 b	264 ab	8.4 a
LSD _{0.05}			23.7	567	28.4

^zMeans for single-plant replicates (n = 5) with 10 subsamples (flowers pollinated) per replicate, except 3x x 3x (n = 3) and 4x x 4x (n = 4).

^yn = total number of fruit harvested per cross.

^xMeans followed by the same letter, within a column, not significantly different based on Fisher's protected LSD at $\alpha = 0.05$.

affect segregation ratios in autotetraploids (Allard, 1960; Ramsey and Schemske, 2002). Random chromatid assortment predicts that crossing-over occurs between the locus and the centromere in tetraploids, which allows for double reduction (sister chromatids in the same gamete). Thus, if we assume random chromatid assortment (quadrivalents with at least one chiasma) we would expect 20.8 non-variegated : 1 variegated progeny in the $F_{2(4x)}$. Our observed ratio was ~10 non-variegated : 1 variegated, falling between the two models. Assuming preferential pairing is limited in recently induced autotetraploids (Wu et al., 2001); the frequency of double reduction would then have a larger impact on allele frequencies and gametic genotypes (Ramsey and Schemske, 2002). In our induced tetraploid $F_{2(4x)}$ s, double reduction would increase the production of homozygous gametes and thus increase the number of nulliplexes (homozygous recessive with four alleles) and the number of variegated progeny. Double reduction depends on the location of a locus on a chromosome relative to the centromere and the rate of chiasma formation, thus double reduction rates will vary among loci. This may explain the increased recovery of variegated progeny and the unaffected frequency of purple progeny if the purple locus is located closer to the centromere. In autotetraploid easter cactus (*Hatiora xgraeseri* Barthlott ex D. Hunt) isozyme segregation ratios fit random chromosomal or chromatidal assortment, depending on the specific loci (Karle et al., 2002), and in allotetraploid quinoa (*Chenopodium quinoa* Willd) discrepancies in tetrasomic segregation ratios for the *R* locus controlling inflorescence color were attributed to variable multivalent formation (Ward, 2000).

Inducing tetraploidy apparently had little effect on male fertility, with pollen staining similar to diploids, however pollen germination slightly decreased compared to that of the diploid. Female fertility, measured as percentage fruit set, was lower in the tetraploids when selfed or crossed with diploids than for diploid selfs. In reciprocal crosses between diploid and tetraploid inbred lines of *Salpiglossis sinuata* R. & P., fruit set did not differ between crosses, however, the number of seeds produced was significantly lower in selfed tetraploids (Needham and Erickson, 1992). For autotetraploids reductions in pollen and seed fertility are usually reduced compared to their diploid progenitor (Ramsey and Schemske, 2002) which is often attributed to the production of multivalents, the frequency of which is negatively correlated with fertility (Sybenga, 1996). In our study, tetraploid selfs had lower percentage fruit set, but average seeds set per fruit and percentage seed germination were not significantly different.

Diploid *H. androsaemum* pollinated with tetraploid pollen ($2x \times 4x$) failed to set fruit. Crosses in this direction commonly fail and violate the endosperm balance number theory, which predicts embryo abortion for crosses that violate a 2 maternal : 1 paternal genomic ratio in developing endosperm (Carputo et al., 1999; Johnston et al., 1980; Sanford, 1983). Crosses with tetraploid *H. androsaemum* as the female ($4x \times 2x$) were successful in producing triploids.

Triploids are expected to be highly infertile owing to uneven segregation from increased multivalent formation and partner exchange during meiosis (Sybenga, 1996) leading to abortive gamete development. Triploids had low pollen germination, even though pollen staining was 40.6%. Pollen staining often results in an overestimation of pollen viability (Olsen et al., 2006) with germination and successful pollination more indicative of viability. Triploid pollen failed to initiate fruit set in selfs and in crosses with tetraploid females in the current study. An additional 139 triploid flowers were selfed on various triploid plants after the

study, resulting in a total of five fruit averaging 41.8 seeds per fruit (data not shown). Diploids pollinated with triploid pollen resulted in just two fruit with reduced seed set and low germination (1%), which may have been the result of diploid pollen contamination or selfing. One fruit was initiated on $3x \times 4x$ crosses, which contained four seeds that failed to germinate. We were unable to recover any viable seeds from triploid female parents regardless of the ploidy level of the male parent. The fertility of triploids are dependent on the production of euploid gametes ($n = x, 2x$, or $3x$) or tolerance of gametes and zygotes to aneuploidy. Triploid blueberries (*Vaccinium corymbosum* L.) are highly sterile, though crosses with tetraploids and hexaploids are variably successful due to the production of $2n$ gametes and tolerance of aneuploid gametes above $2x$ level (Vorsa and Ballington, 1991). Triploid bananas are regarded as sterile, although Ortiz and Vuylsteke (1995) were able to identify clones with residual fertility due to production of euploid gametes from selective elimination of one chromosome set during meiosis. The identification of fertile triploids are valuable for breeding programs; however, selecting infertile triploids are just as valuable for limiting the unwanted spread of ornamental plants in the landscape. The above studies in blueberries and bananas, as well as triploid *Lathyrus pratensis* L. (Khawaja et al., 1997), *S. sinuata* (Needham and Erickson, 1992) and our triploid *H. androsaemum* suggest that hybrid triploids can be screened for infertility as a viable approach for developing new non-invasive cultivars.

Variegated (*var*) and purple (*pl*) foliage in *H. androsaemum* are both simple recessive traits and are not linked, allowing the recovery of the two parental phenotypes as well as the novel combined phenotype (*var/var pl/pl*) at the triploid and tetraploid levels. The greatly reduced male fertility and apparent lack of female fertility among triploids should allow for selection of novel, non-invasive *H. androsaemum* cultivars.

Literature Cited

- Allard, R.W. 1960. Principles of plant breeding. Wiley, New York.
- Alston, F.H., K.L. Phillips, and K.M. Evans. 2000. A *Malus* gene list. Acta Hort. 538:561–570.
- Aluru, M.R. and S.R. Rodermel. 2004. Control of chloroplast redox by the IMMUTANS terminal oxidase. Physiol. Plant. 120:4–11.
- Blinkenberg, C., H. Brix, M. Schaffalitzky de Muckadell, and H. Vedel. 1958. Controlled pollinations in *Fagus*. Silvae Genetica 9:116–122.
- Cadic, A. 1992. Breeding for ever-red barberries (*Berberis* spp.). Acta Hort. 320:85–90.
- Carputo, D., L. Monti, J.E. Werner, and L. Frusciante. 1999. Uses and usefulness of endosperm balance number. Theor. Appl. Genet. 98:478–484.
- Chaparro, J.X., D.J. Werner, R.W. Whetten, and D.M. O'Malley. 1995. Inheritance, genetic interaction, and biochemical characterization of anthocyanin phenotypes in peach. J. Hered. 86:32–38.
- Dias, A.C.P., R.M. Seabra, P.B. Andrade, F. Ferreres, and M. Fernandes-Ferreira. 2000. Xanthone biosynthesis and accumulation in calli and suspended cells of *Hypericum androsaemum*. Plant Sci. 150:93–101.
- Egolf, D.R. 1970. Hibiscus syriacus 'Diana', a new cultivar [Malvaceae]. Bailey 17:75–78.
- Egolf, D.R. 1981. 'Helene' rose of sharon (althea). HortScience 16:226–227.
- Egolf, D.R. 1986. 'Minerva' rose of sharon (althea). HortScience 21:1463–1464.
- Egolf, D.R. 1988. 'Aphrodite' rose of sharon (althea). HortScience 23:223.
- Gustafsson, M.H.G., V. Bittrich, and P.F. Stevens. 2002. Phylogeny of Clusiaceae based on *rbcL* sequences. Intl. J. Plant Sci. 163:1045–1054.

- Hattemer, H.H., W. Steiner, and D. Kownatzki. 1990. Genetic markers in birch. *Silvae Genetica* 39:45–50.
- Heinze, B. and T. Geburek. 1995. Searching for DNA markers linked to leaf colour in copper beech *Fagus sylvatica* L. var. *atropunicea*. *Silvae Genetica* 44:339–343.
- Johnston, S.A., T.P.M. den Nijs, S.J. Peloquin, and R.E. Hanneman, Jr. 1980. The significance of genic balance to endosperm development in interspecific crosses. *Theor. Appl. Genet.* 57:5–9.
- Karle, R., C.A. Parks, M.C. O’Leary, and T.H. Boyle. 2002. Polyploidy-induced changes in the breeding behavior of *Hatiora x graeseri* (Cactaceae). *J. Amer. Soc. Hort. Sci.* 127:397–403.
- Khawaja, H.I.T., J. Sybenga, and J.R. Ellis. 1997. Chromosome pairing and chiasma formation in autopolyploids of different *Lathyrus* species. *Genome* 40:937–944.
- Kihara, H. 1951. Triploid watermelons. *Proc. Amer. Soc. Hort. Sci.* 58:217–230.
- Kirakosyan, A., T.M. Sirvent, D.M. Gibson, and P.B. Kaufman. 2004. The production of hypericins and hyperforin by in vitro cultures of st. john’s wort (*Hypericum perforatum*). *Biotechnol. Appl. Biochem.* 39:71–81.
- Mabberley, D.J. 1997. *The plant book*. Cambridge Univ. Press, Cambridge, U.K.
- Marquard, R.D. 1992. Pollen tube growth in *Carya* and temporal influence of pollen deposition on fertilization success in pecan. *J. Amer. Soc. Hort. Sci.* 117:328–331.
- Matzk, F., A. Meister, and I. Schubert. 2000. An efficient screen for reproductive pathways using mature seeds of monocots and dicots. *Plant J.* 21:97–108.
- McClintock, D., E.C. Nelson, and J. Cassells. 1986. Variegated tutsan, *Hypericum androsaemum* f. *variegatum*. *J. Irish Garden Plant Soc.* 5:25–28.
- McLaren, D.A., E. Bruzzese, and I.G. Pascoe. 1997. The potential fungal pathogens to control *Hypericum* species in Australia. *Plant Protection Qrtly.* 12:81–83.
- Needham, D.C. and H.T. Erickson. 1992. Fecundity of tetraploid x diploid crosses and fertility of the resultant triploids in *Salpiglossis sinuata*. *HortScience* 27:835–837.
- Olsen, R.T., T.G. Ranney, and Z. Vilorio. 2006. Reproductive behavior of induced allotetraploid x *Chitalpa* and in vitro embryo culture of polyploid progeny. *J. Amer. Soc. Hort. Sci.* 131:716–724.
- Ortiz, R. and D. Vuylsteke. 1995. Factors influencing seed set in triploid *Musa* spp. L. and production of euploid hybrids. *Ann. Bot.* 75:151–155.
- Ramsey, J. and D.W. Schemske. 2002. Neopolyploidy in flowering plants. *Annu. Rev. Ecol. Systematics* 33:589–639.
- Ranney, T.G. 2004. Population control: Developing non-invasive nursery crops. *Combined Proc. Intl. Plant Propagators’ Soc.* 54:604–607.
- Reichard, S.H. and P. White. 2001. Horticulture as a pathway of invasive plant introductions in the United States. *BioScience* 51:103–113.
- Robson, N.K.B. 1985. Studies in the genus *Hypericum* L. (Guttiferae). 3. Sections 1. *Campylosporus* to 6a. *Umbraculoides*. *Bul. Brit. Museum (Natural History) Bot. Ser.* 12:163–325.
- Sampson, D.R. and D.F. Cameron. 1965. Inheritance of bronze foliage, extra petals and pendulous habit in ornamental crabapples. *Proc. Amer. Soc. Hort. Sci.* 86:717–722.
- Sanford, J.C. 1983. Ploidy manipulations, p. 100–123. In: J.N. Moore and J. Janick (eds.). *Methods in fruit breeding*. Purdue Univ. Press, West Lafayette, Ind.
- Smith, D.C. and S.A. Mehlenbacher. 1996. Inheritance of contorted growth in hazelnut. *Euphytica* 89:211–213.
- Sybenga, J. 1996. Chromosome pairing affinity and quadrivalent formation in polyploids: Do segmental allopolyploids exist? *Genome* 39:1176–1184.
- Thompson, M.M. 1985. Linkage of the incompatibility locus and red pigmentation genes in hazelnut. *J. Hered.* 76:119–122.
- Tilney-Bassett, R.A.E. 1978. Plastids and variegation: Stable pattern genes, p. 253–285. In: J.T.O. Kirk and R.A.E. Tilney-Bassett (eds.). *The plastids: Their chemistry, structure, growth and inheritance*. Elsevier, New York.
- Valentão, P., E. Fernandes, F. Carvalho, P.B. Andrade, R.M. Seabra, and M.d.L. Bastos. 2002. Antioxidant activity of *Hypericum androsaemum* infusion: Scavenging activity against superoxide radical, hydroxyl radical and hypochlorous acid. *Biol. Pharmaceutical Bul. (Tokyo)* 25:1320–1323.
- Vorsa, N. and J.R. Ballington. 1991. Fertility of triploid highbush blueberry. *J. Amer. Soc. Hort. Sci.* 116:336–341.
- Walbot, V. and E.H. Coe, Jr. 1979. Nuclear gene *iojap* conditions a programmed change to ribosome-less plastids in *Zea mays*. *Proc. Natl. Acad. Sci. USA* 76:2760–2764.
- Ward, S.M. 2000. Allotetraploid segregation for single-gene morphological characters in quinoa (*Chenopodium quinoa* Willd.). *Euphytica* 116:11–16.
- Wu, R., M. Gallo-Meagher, R.C. Littell, and Z.-B. Zeng. 2001. A general polyploid model for analyzing gene segregation in outcrossing tetraploid species. *Genetics* 159:869–882.